

SMALL SEIFERT FIBERED SURGERY ON HYPERBOLIC PRETZEL KNOTS

JEFFREY MEIER

ABSTRACT. We complete the classification of hyperbolic pretzel knots admitting Seifert fibered surgeries. This is the final step in understanding all exceptional surgeries on hyperbolic pretzel knots. We also present results toward similar classifications for non-pretzel Montesinos knots of length three.

1. INTRODUCTION

The study of exceptional surgery on hyperbolic knots has been well developed over the last quarter century. One particularly well studied problem is that of exceptional surgery on arborescent knots, which include Montesinos knots and pretzel knots. Thanks to the positive solution to the Geometrization conjecture [Per03a, Per03b, Per03c], any exceptional surgery is either reducible, toroidal, or a small Seifert fibered space. Exceptional surgeries on hyperbolic arborescent knots of length 4 or greater have been classified [Wu11b], as have exceptional surgeries on hyperbolic 2-bridge knots [BW01]. It has been shown that no hyperbolic arborescent knot admits a reducible surgery [Wu96], and toroidal surgeries on hyperbolic arborescent knots of length 3 are completely classified [Wu11a].

Therefore, it only remains to understand small Seifert fibered surgeries on Montesinos knots of length 3. Furthermore, finite surgeries on Montesinos knots only occur in two instances, along two slopes of each of $P(-2, 3, 7)$ and $P(-2, 3, 9)$ [FIK⁺09, IJ09]. Thus, one must only consider non-finite, atoroidal Seifert fibered surgeries on hyperbolic Montesinos knots of length 3.

According to Wu [Wu10, Wu11c], the only hyperbolic Montesinos knots of length 3 that are pretzel knots and might admit Seifert fibered surgeries have the form $P(q_1, q_2, q_3, n)$, where $n = 0, -1$, $(|q_1|, |q_2|, |q_3|) = (2, |q_2|, |q_3|)$, $(3, 3, |q_3|)$, or $(3, 4, 5)$, and if $n = -1$, then $q_i > 0$ for $i = 1, 2, 3$. Recently, it was shown that hyperbolic pretzel knots of the form $P(p, q, q)$ with p, q positive [IJ11] or $P(-2, p, p)$ with p positive [IJK11] do not admit Seifert fibered surgeries.

Further work by Wu [Wu10, Wu11b, Wu11c] tells us that if a non-pretzel Montesinos knot admits a small Seifert fibered surgery, then it has one of the following forms: $K[1/3, -2/3, 2/5]$, $K[-1/2, 1/3, 2/(2a+1)]$ for $a \in \{3, 4, 5, 6\}$, or $K[-1/2, 1/(2q+1), 2/5]$ for $q \geq 1$.

In this paper, we address the issue of which of the above listed Montesinos knots admit small Seifert fibered surgeries. The main results are stated below. Keep in mind that there is an orientation reversing homeomorphism $K(\alpha) = \overline{K}(-\alpha)$, where $\overline{K}$ is the mirror of K . Thus, we often consider in our analysis, and present in our results, only one representative of $\{K, \overline{K}\}$.

For the following theorem, recall that the pretzel knot $P(p, q, r)$ with $|p|, |q|, |r| \geq 2$ is hyperbolic unless it is either $P(-2, 3, 3)$ or $P(-2, 3, 5)$, in which case it is the torus knot $T(3, 4)$ or $T(3, 5)$, respectively ([Oer84]). Below, when we consider the knots $P(-2, 2p +$

$1, 2q + 1)$, we will assume that $|p| < |q|$ when p and q have the same sign and that $p > 0$ when their signs differ.

Theorem 1.1. The hyperbolic pretzel knot $P(-2, 2p + 1, 2q + 1)$, with the conventions discussed above, admits a small Seifert fibered surgery if and only if $p = 1$, in which case it admits precisely the following small Seifert fibered surgeries:

- $P(-2, 3, 2q + 1)(4q + 6) = S^2(1/2, -1/4, 2/(2q - 5))$
- $P(-2, 3, 2q + 1)(4q + 7) = S^2(2/3, -2/5, 1/(q - 2))$

Theorem 1.2. Hyperbolic pretzel knots of the form $P(3, 3, m)$ or $P(3, 3, m, -1)$ admit no small Seifert fibered surgeries. Hyperbolic pretzel knots of the form $P(3, -3, m)$, with $m < 0$, admit small Seifert fibered surgeries precisely in the following cases:

- $P(3, -3, -2)(-1) = S^2(2/3, -1/4, -2/5)$
- $P(3, -3, -3)(-1) = S^2(-1/2, 1/5, 2/7)$
- $P(3, -3, -4)(-1) = S^2(1/2, -1/5, -2/7)$
- $P(3, -3, -5)(-1) = S^2(1/3, 1/4, -3/5)$
- $P(3, -3, -6)(-1) = S^2(1/2, -1/3, -2/13)$

Theorem 1.3. Pretzel knots of the form $P(3, \pm 4, \pm 5)$ admit no small Seifert fibered surgeries.

Theorem 1.4. Suppose that K is a non-pretzel Montesinos knot and $K(\alpha)$ is a small Seifert fibered space. Then either $K = K[-1/2, 2/5, 1/(2q + 1)]$ for some $q \geq 5$, or K is on the following list and has the described surgeries.

- $K[1/3, -2/3, 2/5](-5) = S^2(1/4, 2/5, -3/5)$
- $K[-1/2, 1/3, 2/7](-1) = S^2(1/3, 1/4, -4/7)$
- $K[-1/2, 1/3, 2/7](0) = S^2(1/2, 3/10, -4/5)$
- $K[-1/2, 1/3, 2/7](1) = S^2(1/2, 1/3, -16/19)$
- $K[-1/2, 1/3, 2/9](2) = S^2(1/2, -1/3, -3/20)$
- $K[-1/2, 1/3, 2/9](3) = S^2(1/2, -1/5, -3/11)$
- $K[-1/2, 1/3, 2/9](4) = S^2(-1/4, 2/3, -3/8)$
- $K[-1/2, 1/3, 2/11](-2) = S^2(-2/3, 2/5, 2/7)$
- $K[-1/2, 1/3, 2/11](-1) = S^2(-1/2, -2/7, 2/9)$
- $K[-1/2, 1/3, 2/5](3) = S^2(1/2, -1/3, -2/15)$
- $K[-1/2, 1/3, 2/5](4) = S^2(1/2, -1/6, -2/7)$
- $K[-1/2, 1/3, 2/5](5) = S^2(-1/3, -1/5, 3/5)$
- $K[-1/2, 1/5, 2/5](7) = S^2(1/2, -1/5, -2/9)$
- $K[-1/2, 1/5, 2/5](8) = S^2(-1/4, 3/4, -2/5)$
- $K[-1/2, 1/7, 2/5](11) = S^2(-1/3, 3/4, -2/7)$

Each of the theorems stated above are proved below using a common procedure. First, we exploit the symmetries of the Montesinos knots in question to describe the surgery space as a branched double cover of a link. Next, we use rational tangle filling theory and exceptional surgery bounds to restrict our attention to a finite list of such links, i.e, we restrict the parameters for which the Montesinos knots in question can admit small Seifert fibered surgeries. Finally, we use knot theory invariants to show that the branched double covers of links on this finite list cannot be Seifert fibered (excepting, of course, the cases that are). This last step makes use of the Mathematica[®] package KnotTheory[‘] [Wol99].

It should be noted that, concurrent with the preparation of this paper, the author learned that similar results had been obtained by Wu, though using different techniques. Wu also restricted the families to finite families of surgery spaces, but did so by studying exceptional surgery on *tubed* Montesinos knots (see [Wu12a]). He then appealed to a computer program called *Snappex*, to determine the hyperbolic structure of the surgeries in question (see [Wu12b]). At the time of writing, it was not clear whether or not Wu's program had successfully checked each of the remaining cases.

1.1. A word on non-integral surgeries. In a survey by Wu [Wu98a], it is shown how techniques and results from [Bri98, Wu98b] can be combined with work of Delman [Del95] to study which length-3 Montesinos knots have exteriors that admit persistent essential laminations.

Theorem 1.5. Let K be a hyperbolic Montesinos knot of length 3. Then the exterior of K admits a persistent essential lamination, and, thus, cannot admit a non-integral small Seifert fibered surgery, unless $K = K[x, 1/p, 1/q]$ (or its mirror image), where $x \in \{-1/(2n), -1 \pm 1/(2n), -2 + 1/(2n)\}$, and p, q , and n are positive integers.

With this in mind, for many of the families of pretzel knots considered in this paper, it is only necessary to consider integral surgeries. However, for some families, it is necessary to consider non-integral surgeries. To be specific, of all the pretzel knots considered in this paper, only the following families could potentially admit non-integral small Seifert fibered surgeries:

- $P(-2, 2p + 1, 2q + 1)$ with $1 \leq p < q$
- $P(3, 3, -2m)$ with $m \geq 2$
- $P(3, 3, 2m, -1)$ with $m \geq 1$
- $P(3, -4, 5)$

Thus, whenever such a family is considered, we have shown that, in fact, there are no non-integral small Seifert fibered surgeries. One of the biggest open problems in the study of exceptional Dehn surgery is the following conjecture (see [Gor09]).

Conjecture 1.6. Any Seifert fibered surgery on a hyperbolic knot is integral.

The results of this paper are the final steps of an affirmative answer to Conjecture 1.6 in the case of hyperbolic arborescent knots.

Theorem 1.7. Any Seifert fibered surgery on a hyperbolic arborescent knot is integral.

1.2. Open questions. Unfortunately, the techniques of this paper are insufficient to complete the classification of Seifert fibered surgery on Montesinos knots. We are left with the following question, which is the final step in a complete classification of exceptional surgery on arborescent knots.

Question 1.8. Do the Montesinos knots $K[-1/2, 2/5, 1/(2q + 1)]$ with $q \geq 5$ admit small Seifert fibered surgeries?

2. PRELIMINARIES

2.1. Dehn surgery. Let K be a knot in S^3 , and let $N(K)$ be a regular neighborhood of K . Let $M_K = \overline{S^3 \setminus N(K)}$ be the exterior of K . The set of isotopy classes of simple closed curves on $\partial N(K) = \partial M_K$ is in bijection with $H_1(\partial M_K)$, the latter of which is naturally generated




FIGURE 2.1. On the right, we see the exterior, M_K , of the left-handed trefoil. The surgery space $K(0)$ is formed by filling the boundary of M_K with a solid torus such that the meridian maps to a 0-slope (a longitude of K) on ∂M_K .

by two elements: $[\mu]$ and $[\lambda]$, where $[\mu]$ generates $H_1(M_K) \cong \mathbb{Z}$, $[\lambda] = 0 \in H_1(M_K)$, and μ and λ intersect geometrically once on ∂M_K . Orient μ and λ so that $\mu \cdot \lambda = +1$. The unoriented isotopy class of a simple closed curve $\gamma \subset \partial M_K$ is called a *slope* and can be thought of as an element $m/l \in \mathbb{Q} \cup \{\infty\}$, where $[\gamma] = m[\mu] + l[\lambda]$ in $H_1(\partial M_K)$. The curves μ and λ are called the *meridian* and the *longitude*, respectively.

Given two slopes α and β on T^2 , let the *distance* between α and β , $\Delta(\alpha, \beta)$ be their minimal geometric intersection number. If $\alpha = m/l$ and $\beta = m'/l'$, then we have $\Delta(\alpha, \beta) = |ml' - m'l|$.

Let V be a solid torus, and let $\varphi : \partial V \rightarrow \partial M_K$ be a homeomorphism which takes the meridian of V to a slope α on ∂M_K . Then *Dehn surgery on K along α* , or α -*Dehn surgery on K* , is the space $K(\alpha) = M_K \cup \varphi V$. See Figure 2.1. For a general overview of the theory of Dehn surgery, a subject that has been well-studied since its introduction by Dehn in 1910 [Deh10], see [Gor09].

Dehn surgery generalizes nicely to manifolds M with a torus boundary component $T \subset \partial M$, where M may not be the complement of knot in S^3 . Let $\alpha \subset T$ be a slope, then α -*Dehn filling of M on T* is the space $M(\alpha) = M \cup_\varphi V$, where $\varphi : \partial V \rightarrow T$ sends the meridian of V onto α . One difference in this scenario is that there may be no canonical way to distinguish a longitude on T , however, $\Delta(\alpha, \beta)$ is still well-defined for any pair of slopes, α and β .

2.2. Cable spaces. Let V be a solid torus, and let J be a (p, q) -curve inside V (see Figure 2.2). The *cable space*, $C(p, q)$, is the space formed by removing a regular neighborhood of J . Let $T_1 = \partial V$ and $T_0 = \partial N(J)$. There is a properly embedded annulus, A , connecting the two boundary components such that $A \cap T_1$ is a p/q -curve (in terms of the standard meridian and longitude on V) and $\gamma = A \cap T_0$ is a $pq/1$ -curve (see Figure 2.2). Let μ and λ be some choice of meridian and longitude for T_0 . Then the slope γ is called the *cabling slope* for $C(p, q)$.

Let A' be a properly embedded annulus such that $A' \cap T_0$ is two $pq/1$ -curves, parallel to each other and to γ (see Figure 2.2). Now, let $C(p, q)(\alpha)$ denote α -Dehn filling on T_0 . Then, if $\alpha = \gamma$, this filling has the effect of capping off A' to form a separating 2-sphere, S , and capping off one boundary component of A to form a disk, D , which intersects T_1 in a p/q -curve. The result is that $C(p, q)(\gamma) = (S^1 \times D^2) \# L(q, p)$.




FIGURE 2.2. On the left, we see a $(4, 5)$ -curve J inside a solid torus, V , and, on the right, we see a cross section of $V - N(J)$, along with two interesting annuli, A and A' .

Let $t : C(p, q) \rightarrow C(p, q)$ represent Dehn twisting along A . Then, $t^l(\mu) = \mu + l(pq\mu + \lambda) = (lpq + 1)\mu + l\lambda$. Since $C(p, q)(\mu) = S^1 \times D^2$, it follows that $C(p, q)(t^l(\mu)) = S^1 \times D^2$. So, slopes of the form $(lpq + 1)/l$ all correspond to surgery slopes on T_0 that yield solid tori.

This shows that cable spaces have infinitely many fillings returning solid tori, all at distance one from the cabling slope.

On the other hand, we have the following lemma, which follows from the Cyclic Surgery Theorem [CGLS87] and work of Gabai [Gab89]. See [Kan10] for a proof and more general discussion.

- Lemma 2.1.** (a) Let $M \neq T^2 \times I$ be an irreducible and ∂ -irreducible 3-manifold with a torus boundary component, T_0 . Let T_1 be an incompressible torus in M , distinct from T_0 . If α and β are slopes on $T_0 \subset \partial M$ with $\Delta(\alpha, \beta) \geq 2$, such that T_1 is compressible in $M(\alpha)$ and $M(\beta)$, then M is a cable space with cabling slope γ such that $\Delta(\alpha, \gamma) = \Delta(\beta, \gamma) = 1$.
- (b) Let $M \neq T^2 \times I$ be an irreducible and ∂ -irreducible 3-manifold with a torus boundary component, T_0 . Let T_1 be an incompressible torus in M , distinct from T_0 . If α and β are slopes on $T_0 \subset \partial M$ with $\Delta(\alpha, \beta) = 1$, such that T_1 is compressible in $M(\alpha)$ and $M(\beta)$, then either
- (a) M is a cable space with cabling slope α or β , or
 - (b) M is the exterior of a braid in a solid torus, $M(\alpha)$ and $M(\beta)$ are solid tori, and $\Delta(\eta_\alpha, \eta_\beta) \geq 4$, where η_α and η_β are the induced slopes of the meridian on T_0 .

2.3. Seifert fibered spaces. A *fibered solid torus* is formed by gluing the ends of $D^2 \times I$ together with a twist ρ through $\frac{2\pi p}{q}$, where $q \geq 1$ and p and q are relatively prime. There are two types of *fibers*: the *central fiber*, i.e., the image of $(0, 0) \times I$ after gluing, and the union of the arcs $x \times I, \rho(x) \times I, \dots, \rho^{q-1}(x) \times I$, for $x \neq (0, 0)$.

A *Seifert fibered space* is a 3-manifold that can be decomposed as a disjoint union of circles (called *fibers*), where each fiber has a regular neighborhood homeomorphic to a fibered solid torus, i.e., the fiber becomes the central fiber of the fibered solid torus. Viewing the neighborhood this way, if $q = 1$, we say the fiber is *ordinary*. If $q \geq 2$, we say the fiber is *exceptional with multiplicity q* . In the latter case, the fibers surrounding the central fiber are called (p, q) -curves.

If M is a Seifert fibered space, there is a natural projection $\pi : M \rightarrow \Sigma$ that identifies each fiber to a point. The surface Σ is called the base space. We can record the exceptional

fiber information in the form of cone points on Σ , so M is a circle bundle over the resulting orbifold. Another way to recover M is to remove a disk neighborhood of each cone point on Σ and cross the resulting surface with S^1 . The result is a manifold with torus boundary components. If we choose meridian and longitude coordinates for each boundary component so that the projection of the meridians to the base surfaces is one-to-one onto the boundary of the removed disks and the longitude is $\star \times S^1$ in the circle product, then M is the result of Dehn filling on the boundary components along the slopes p'/q , where $pp' \equiv 1 \pmod{q}$. If M is a Seifert fibered space with base space Σ and n exceptional fibers with fibered solid torus neighborhoods consisting of (p_i/q_i) -curves for $i = 1, 2, \dots, n$, we write $M = \Sigma(p'_1/q_1, \dots, p'_n/q_n)$, or sometimes $M = \Sigma(q_1, \dots, q_n)$. In fact, the homeomorphism type of M is determined by Σ and the *Seifert invariants*: $\{p'_1/q_1, \dots, p'_n/q_n\}$, up to permutation, and up to the relation $\{p'_1/q_1, p'_2/q_2, \dots, p'_n/q_n\} = \{p'_1/q_1 \pm 1, p'_2/q_2 \mp 1, \dots, p'_n/q_n\}$. In other words, $\sum_{i=1}^n p'_i/q_i$ is an invariant of M . Because of this, it is often useful to standardize the notation so that the Seifert invariants are all positive and less than one. To do this, we subtract out the integer part of each fraction and collect them in a single term, b . We write $M = \Sigma(b; p'_1/q_1, \dots, p'_n/q_n)$, where $0 < p'_i < q_i$ and $b \in \mathbb{Z}$.

A Seifert fibered space is called *small* if the base space is a sphere and the number of exceptional fibers is at most three.

Next, we recall a fact about Dehn filling on Seifert fibered manifolds that will be useful throughout this paper. Let M be a Seifert fibered manifold with a torus boundary component $T \subset \partial M$. The fibering of M induces a fibering of T , and the slope, γ , of the induced fibers on T is called the *Seifert slope* of T . Now, the Seifert fibering of M will extend to a Seifert fibering of α -Dehn filling on M provided that $\alpha \neq \gamma$. In fact, we have the following. See [Hei74] for a complete treatment of Dehn filling on Seifert fibered spaces with boundary.

Lemma 2.2. If M is a Seifert fibered manifold with base surface Σ and n exceptional fibers, $T \subset \partial M$ is a torus boundary component (corresponding to a circle boundary component $C \subset \partial \Sigma$), and γ is the Seifert slope of T , then let $M(\alpha)$ denote α -Dehn filling on T , let $d = \Delta(\alpha, \gamma)$, and let $\hat{\Sigma} = \Sigma \cup_C D^2$. Then,

- (a) If $d \geq 2$, $M(\alpha)$ is a Seifert fibered space with base surface $\hat{\Sigma}$ and $n+1$ exceptional fibers (the original exceptional fibers, plus a new one of multiplicity d).
- (b) If $d = 1$, $M(\alpha)$ is a Seifert fibered space with base surface $\hat{\Sigma}$ and (the original) n exceptional fibers.
- (c) If $d = 0$, $M(\alpha) = N \# L$, where N is a Seifert fibered space with base surface $\hat{\Sigma}$ and (the original) n exceptional fibers, and L is a Lens space.

As an example, consider $D^2(a, b)$ with Seifert slope r/s , and let $d = \Delta(m/l, r/s) = |ms - lr|$. Then (as developed in [Gor09]),

$$D^2(a, b)(m/l) = \begin{cases} S^2(a, b, d) & \text{if } d \geq 2 \\ L(m, lb^2) & \text{if } d = 1 \\ L(a, b) \# L(b, a) & \text{if } d = 0 \end{cases}$$

2.4. Montesinos knots. A *tangle* is a pair (B, A) , where $B \cong B^3$ and A is a pair of properly embedded arcs in B . A *marked tangle* is a tangle along with an identification of its boundary $\partial(B, A) = (S, S \cap A)$, which is a 2-sphere with 4 distinguished points, with the pair $(S^2, \{NE, NW, SW, SE\})$. The *trivial tangle* is the tangle which is homeomorphic as a




FIGURE 2.3. Above we have the Montesinos knot $K[1/3, 1/4, -3/5]$ and the 2-bridge knot $K[43/95]$ (with continued fraction $[2, 4, 1, 3, 2]$).

marked tangle to $(D^2, \{2 \text{ points}\}) \times I$. Let h and r be the tangle operations where h adds a positive horizontal half-twist (right-handed), and r is reflection in the (NW/SE) -plane.

Let $[c_1, c_2, \dots, c_m]$ be a sequence of integers, and let $p/q = \frac{1}{c_1 + \frac{1}{c_2 + \frac{1}{\dots + \frac{1}{c_m}}}}$. The *rational tangle*, $\mathcal{R}(p/q)$ is formed by applying the operation $(h^{c_m}r)(h^{c_{m-1}}r) \dots (h^{c_1}r)$ to the trivial tangle, which we denote $\mathcal{R}(1/0)$. Note that, as an unmarked tangle, $\mathcal{R}(p/q)$ is trivial, one can just untwist it. On the other hand, Conway showed [Con70] that, as marked tangles, $\mathcal{R}(p/q) = \mathcal{R}(p'/q')$ if and only if $p/q = p'/q'$.

A *Montesinos link of length n* is a link formed by connecting n rational tangles to each other in a standard fashion. We denote such a knot by $K(p_1/q_1, \dots, p_n/q_n)$ (see Figure 2.4). In the special case where each $p_i = \pm 1$, we have what is called a *pretzel knot*. In this case, each tangle is just a strand of vertical twists, since $1/q$ has the continued fraction expansion $[q]$. It is easy to see that Montesinos links of length one or two are the same. These links are called *2-bridge links*, and will be denoted $K[p/q]$, where p/q is the rational number describing the tangle twists.

Montesinos links are determined up to the same relations as Seifert fibered spaces, so we can normalize the invariants and write $K(b; p_1/q_1, \dots, p_n/q_n)$ where $0 < p_i < q_i$ and $b \in \mathbb{Z}$. In fact, we have the following proposition, which follows from Theorem 2.5 below.

Proposition 2.3. The double cover of S^3 , branched along the Montesinos link $K[p_1/q_1, \dots, p_n/q_n]$, is the Seifert fibered space $S^2(p_1/q_1, \dots, p_n/q_n)$.

We remark that it is often helpful to allow p_i/q_i to be zero, ∞ , or 1 for some i , in either the notation for Montesinos links or Seifert fibered spaces. For our purposes, this will only happen when the length n is three or less, and the result should be clear from the context. For example, $K[1/3, -1/2, 1/0]$ is the connected sum of a trefoil knot and a Hopf link, $K[1/3, 2/7, 0] = K[2/13]$, and $S^2(2, 3, 1)$ is a lens space.

2.5. Seifert fibered surgery on knots with symmetries. In this section, we recall some known results about Seifert fibered surgery on knots that admit a strong inversion, have period two, or both. In what follows, let $K \subset S^3$ be a knot and let $\varphi : S^3 \rightarrow S^3$ be a non-trivial orientation preserving involution such that $\varphi(K) = K$ and $C_\varphi = \text{Fix}(\varphi) \neq \emptyset$. By the positive solution to the Smith conjecture, C_φ is an unknotted circle in S^3 [MB84].

Definition 2.4. If $C_\varphi \cap K \neq \emptyset$, then φ is called a *strong inversion* of K and K is called *strongly invertible*. In this case, $C_\varphi \cap K = 2$ points and φ reverses the orientation of K .

If $C_\varphi \cap K = \emptyset$, then we say φ is a *cycle of order 2* and that K has *period 2*.




FIGURE 2.4. The knot $P(3, 3, -6)$, shown with its three symmetries and the resulting quotients.

In this paper, we will only be interested in strong inversions and cycles of period 2. For a more general treatment of Dehn surgery on knots with symmetries, see [Mot03].

First, let us consider strongly invertible knots. Let $K \subset S^3$ be a knot with a strong inversion φ . Then φ restricts to an involution of the knot exterior, M_K , and the quotient of M_K by the action of φ is a tangle, $\mathcal{T}_K$. The well-known Montesinos trick gives a correspondence between Dehn filling on M_K and rational tangle filling on $\mathcal{T}_K$. For details, see [Gor09]. The following is originally due to Montesinos [Mon75].

Theorem 2.5. Let $\mathcal{T}$ be a marked tangle. Then $\widetilde{\mathcal{T}}(r/s) \cong \widetilde{\mathcal{T}}(-r/s)$.

Let $L_{r/s} = \mathcal{T}_K(-r/s)$, so $L_{r/s}$ is a knot or a two-component link in S^3 with $K(r/s)$ as the double cover of S^3 , branched along $L_{r/s}$. Suppose that $K(r/s)$ is a small Seifert fibered space. Let $\bar{\varphi} : K(r/s) \rightarrow K(r/s)$ be the involution induced by extending φ across the filling solid torus. Then $K(r/s)/\bar{\varphi} = S^3$.

If K is not a trefoil knot, then we can assume that $\bar{\varphi}$ is fiber-preserving [Mot03]. Let $\pi : K(r/s) \rightarrow S^2$ be the Seifert fibration of $K(r/s)$. Let $C_{\bar{\varphi}} = \text{Fix}(\bar{\varphi})$. If each component of $C_{\bar{\varphi}}$ is a fiber in $K(r/s)$, then $K(r/s) \setminus C_{\bar{\varphi}}$ admits a Seifert fibered structure. Since this structure is compatible with $\bar{\varphi}$, $S^3 \setminus L_{r/s}$ admits a Seifert fibered structure. In other words, $L_{r/s}$ is a *Seifert link*.

Let $\hat{\varphi} : S^2 \rightarrow S^2$ be the induced involution of the base orbifold. If one component of $C_{\bar{\varphi}}$ is not a fiber in $K(r/s)$, then $\hat{\varphi}$ is reflection across the equatorial circle, $C_{\hat{\varphi}}$, of S^2 and all of the cone points lie on $C_{\hat{\varphi}}$ [Mot03]. In this case, $L_{r/s} = C_{\bar{\varphi}}/\bar{\varphi}$ is a length-3 Montesinos link [MM02]. So, we have the following, as stated in [IJ11].

Proposition 2.6. Let K be a strongly invertible hyperbolic knot, and let $r/s \in \mathbb{Q}$. Let $L_{r/s}$ be the link obtained by applying the Montesinos trick to $K(r/s)$. If $K(r/s)$ is a small




FIGURE 2.5. Possible configurations of cone points in the base sphere of a small Seifert fibered space

Seifert fibered space with base orbifold S^2 , then $L_{r/s}$ is either a Seifert link or a Montesinos link.

Seifert links are well understood [BM70, EN85]. In the present paper, we will only be concerned with Seifert knots and Seifert links with two components, in which case we have the following.

Lemma 2.7. Let $L \subset S^3$ be a Seifert link with at most two components. Then L is equivalent to one of the following:

- (a) A torus knot
- (b) A two-component torus link
- (c) A two-component link consisting of a torus knot together with a core curve of the torus on which it lies.

Note that, in particular, every component of a Seifert link is a torus knot or an unknot.

Now let $K \subset S^3$ be a knot with a cycle φ of order 2. Suppose that $K(r/s)$ is a Seifert fibered space with base surface S^2 , and let $\bar{\varphi}$ be the extension of $\varphi|_{M_K}$ to $K(r/s)$. Then, $K(r/s)$ has a $\bar{\varphi}$ -invariant Seifert fibered structure [MM02]. Let $C_{\bar{\varphi}} = \text{Fix}(\bar{\varphi})$, and let $L_{r/s} = C_{\bar{\varphi}}/\bar{\varphi}$.

If r is odd, then $L_{r/s}$ is a knot. If r is even, then $L_{r/s}$ is a link. Let $K_{\varphi} = K/\varphi$. K_{φ} is called the *factor knot* of K (with respect to φ), and let $C_{\varphi} = \text{Fix}(\varphi)$. In the case where r is odd, we can view $L_{r/s}$ as the image of C_{φ}/φ after $r/2s$ surgery on K_{φ} , so $L_{r/s}$ is a knot in $K_{\varphi}(r/2s)$. If r is even, then $L_{r/s}$ is the image of C_{φ}/φ in $K_{\varphi}(r/2s)$ together with the core of the surgery torus, so $L_{r/s}$ is a link in $K_{\varphi}(r/2s)$.

Let $\pi : K(r/s) \rightarrow S^2$ be a Seifert fibering of $K(r/s)$, and let $\hat{\varphi}$ be the induced involution of S^2 , with fixed point set $C_{\hat{\varphi}}$. In [MM02], it is shown that if K is not a torus knot or a cable of a torus knot, then no component of $C_{\bar{\varphi}}$ is a fiber in $K(r/s)$ and $C_{\hat{\varphi}}$ is the equatorial circle in S^2 . This implies that $\hat{\varphi}$ is reflection across the equator. Since $\bar{\varphi}$ is fiber preserving, $\hat{\varphi}$ must pair up cone points in the northern hemisphere with cone points in the southern hemisphere. Let k denote the number of cone points in the northern hemisphere. For our purposes, $k = 0$ or $k = 1$. From [MM02], we have the following:

Lemma 2.8. (1) If $k = 0$, then $K_{\varphi}(r/2s) = K(r/s)/\bar{\varphi} \cong S^3$.
 (2) If $k = 1$, then $K_{\varphi}(r/2s) = K(r/s)/\bar{\varphi}$ is a lens space..

Note that S^3 and $S^2 \times S^1$ are not lens spaces. These facts can be helpful in obstructing Seifert fibered surgeries, based on the knot type of K_φ . Throughout, U will represent the unknot.

Corollary 2.9. Let $K \subset S^3$ be a period 2 hyperbolic knot with factor knot K_φ . Suppose $K(r/s)$ is a small Seifert fibered space.

- (1) If $K_\varphi = T_{p,q}$ and r is even, then $\Delta(pq, r/2s) = 1$, so $|r - 2spq| = 2$.
- (2) If $K_\varphi = T_{p,q}$ and r is odd, then $\Delta(pq, r/2s) = 1$, so $|r - 2spq| = 1$.
- (3) If $K_\varphi = U$ and $k = 0$, then $|r| \leq 2$.
- (4) If $K_\varphi = U$ and $k = 1$, then $|r| \geq 3$.

Proof. If $K(r/s)$ is a small Seifert fibered space, then $K_\varphi(r/2s)$ is a lens space if $k = 1$ and S^3 if $k = 0$. Such surgeries on U and $T_{p,q}$ are well understood (see [Gor09]). $\square$

2.6. Some exceptional Dehn surgery results. There are many important results in the study of exceptional Dehn surgery that give limitations on which slopes can be exceptional for a hyperbolic knot K . Below, we present some of the results that will be used in this paper. First, we state an important result of Lackenby and Meyerhoff [LM08] that tells us that exceptional fillings are always “close” to each other.

Theorem 2.10. Suppose M is a hyperbolic manifold with torus boundary component $T \subset \partial M$ and that α and β are exceptional filling slopes on T . Then $\Delta(\alpha, \beta) \leq 8$.

The distance bound of 8 above can be improved if one specifies the type of space for each of $M(\alpha)$ and $M(\beta)$. Let S and T represent the sets of reducible and toroidal manifolds, respectively. Let L represent the set of lens spaces. Let $\Delta(C_1, C_2)$ represent the largest possible value of $\Delta(\alpha, \beta)$ such that there exists a hyperbolic manifold M with $M(\alpha)$ a manifold of type C_1 and $M(\beta)$ a manifold of type C_2 (we will always consider manifolds with one boundary component, though the theory is more general). The following table presents the known values of $\Delta(C_1, C_2)$.

	S	T	S^3	L
S	1	3	?	1
T		8	2	?
S^3			0	1
L				1

Notice that in the case of (S, S^3) , this is equivalent to the cabling conjecture, and in the case of (T, L) , the bound is known to be either 3 or 4 [Lee11]. For a more thorough discussion of these bounds, the manifolds achieving them, and precise references, see [Gor09] and [Gor99].

Suspiciously absent from the above table are bounds on the distance between a (non-lens space) small Seifert fibered surgery and the other types of exceptional surgeries. These seem to be the most difficult cases to analyze, and, in particular, it is not known whether or not the distance 8 bound of Lackenby and Meyerhoff can be improved in most of the cases (though, see Theorems 2.11 and 2.12 below).

Recent work in progress by Boyer, Gordon, and Zhang [BGZ12] gives a stronger bound on the distance between toroidal and small Seifert fibered fillings. The proof is complicated, and not all cases have yet been accounted for. However, we have the following, which can be seen by piecing together Corollary 7.6, Proposition 14.1, and Proposition 16.1 in [BGZ12].

Theorem 2.11. For any hyperbolic manifold M , if $M(\alpha) = A \cup_{T^2} B$ is toroidal with one of A or B non-Seifert fibered, then for any slope β such that $M(\beta)$ is a Seifert fibered space, $\Delta(\alpha, \beta) \leq 6$.

Since many of the pretzel knots studied below are genus one, it will be helpful for us to have the following result, which gives particularly strong bounds on small Seifert fibered surgery slopes [BGZ11] of such knots.

Theorem 2.12. Let M be a hyperbolic manifold with boundary component $T \subset \partial M$. Suppose that M contains a properly embedded essential once-punctured torus F such that $F \cap T$ has slope α . If $M(\alpha)$ is a small Seifert fibered space, then $\Delta(\alpha, 0) \leq 3$.

2.7. Invariants from knot theory and Montesinos and torus links. In this section we give a very brief overview of some knot and link invariants and how they will be used to obstruct the quotient links encountered in this paper from being Seifert links or Montesinos links. We will present a series of criteria that will be applied in each of the following sections.

A link is called *k-almost alternating* if it has a k -almost alternating diagram, but no $(k-1)$ -almost alternating diagram, i.e., if it has a diagram D such that changing k crossings of D gives a new diagram that is alternating, but no such diagram where the same result is achieved after $k-1$ crossing changes.

Recall that the Khovanov homology, $\text{Kh}(L)$, is a bi-graded abelian group associated to L , and that the *width* of $\text{Kh}(L)$ is the number of diagonals that support a nontrivial element in $\text{Kh}(L)$. Denote this width by $|\text{Kh}(L)|$. Then we have the following theorem. (See, for example, [AP04].)

Theorem 2.13. Let L be a non-split k -almost alternating link. Then $|\text{Kh}(L)| \leq k + 2$.

An easy exercise shows that any length-3 Montesinos link is almost alternating (i.e., 1-almost alternating), so we have our first obstruction criterion.

Criterion 2.14. If $|\text{Kh}(L)| \geq 4$, then L is not a length-3 Montesinos link.

When we encounter links L that do not satisfy this criterion, then we will use the following program to show they are not a Montesinos link. We will generate a list of all Montesinos links whose crossing numbers are compatible with that of L (i.e., less than the number of crossings in a diagram of L). (Note that the crossing number of a Montesinos link is well understood [LT88].) We will then check this list for elements that, if L is a knot, have the same determinant, Alexander polynomial, Jones polynomial, Khovanov homology, and, if need be, Kauffman polynomial or HOMFLYPT polynomial, and that, if L is a 2-component link, have the same determinant, Jones polynomial, Khovanov homology, and, if need be, Kauffman polynomial or HOMFLYPT polynomial. We will refer to this method as *Method 1*. This very large number of computations was performed using the KnotTheory[®] package for Mathematica[®] [Wol99].

Next, we observe that if K is a length-3 Montesinos link, then it is the union of 2-bridge knots and unknots. If K is the union of two unknots, then it has the form $K[p_1/q_1, p_2/q_2, p_3/q_3]$, where each p_i is even. If one component of K is the 2-bridge knot $K[p/q]$, then K has the form $K[p_1/q_1, p_2/q_2, x/q]$ with q_1 and q_2 even and with $x = p$ or $\bar{p}$, where $p\bar{p} \equiv 1 \pmod{q}$. If we consider the unknot a 2-bridge knot, then we have the following criterion.

Criterion 2.15. If K is a 2-component link such that one component is not a 2-bridge knot, then K is not a Montesinos link.

Using Method 1 and Criteria 2.14 and 2.15, any knot or link we encounter that we claim is not a Montesinos knot or link can be shown to not be a Montesinos knot or link.

Now we recall some facts about torus knots (see, for example, [Cro04]). Let $T(p, q)$ be the (p, q) -torus link for $p, q \geq 2$, where $T(p, q)$ is a knot if and only if p and q are coprime. Then, $T(p, q)$ is a positive link, i.e., has a diagram with all positive crossings. Furthermore, in the case of a torus knot, $2g(T(p, q)) = (p - 1)(q - 1)$, where $g(K)$ denotes the genus of the knot K , and $\det(T(p, q)) = p$ if q is even, and 1 if both p and q are odd, where $\det(L)$ denotes the determinant of the link L . Let $s(K)$ denote the Rasmussen invariant of K , as defined in [Ras10], where the following was shown.

Proposition 2.16. If K is a positive knot, then $s(K) = 2g(K)$.

Recall that $2g(K)$ is bounded below by the breadth of the Alexander polynomial, which we denote $\text{br}(\Delta_K(t))$. This gives us the following criterion.

Criterion 2.17. If $s(K) < \text{br}(\Delta_K(t))$ or $2g(K) \neq s(K)$, then K is not a torus knot.

We also have, by our discussion above:

Criterion 2.18. If $\det(K) > s(K) + 1$, then K is not a torus knot.

If we consider the unknot a torus knot, then each component of a two component Seifert link is a torus knot, so we have the following criterion.

Criterion 2.19. If L is a two-component link such that a component is not a torus knot, then K is not a Seifert link.

In what follows, Criteria 2.17, 2.18, and 2.19 often suffice to prove that a link is not a Seifert link. In the few cases where they fail, further argument is given to accomplish the feat.

3. THE CASE OF $(2, |q_2|, |q_3|)$

Let $K_{p,q}$ be the hyperbolic pretzel knot $P(-2, 2p+1, 2q+1)$ (see Figure 3.1). Since Ichihara and Jong have shown that $K_{p,p}$ admits no small Seifert fibered surgery [IJ11, IJK11], and by interchanging p and q , if necessary, we may assume that $|q| > |p|$ if p and q have the same sign and $p > 0$ otherwise. Let $\alpha_r = 4(p + q + 1) - r$, for $r \in \mathbb{Q}$.

Note that α_r is chosen this way so that α_r -surgery on $K_{p,q}$ will correspond with r -filling of $\mathcal{T}_{p,q}$. This becomes clear if one carefully follows through the process of obtaining $\mathcal{T}_{p,q}$ from $K_{p,q}$ by applying the Montesinos trick and isotoping.

Our first result is the following.

Theorem 1.1. The hyperbolic pretzel knot $P(-2, 2p + 1, 2q + 1)$, with the conventions discussed above, admits a small Seifert fibered surgery if and only if $p = 1$, in which case it admits precisely the following small Seifert fibered surgeries:

- $P(-2, 3, 2q + 1)(4q + 6) = S^2(1/2, -1/4, 2/(2q - 5))$
- $P(-2, 3, 2q + 1)(4q + 7) = S^2(2/3, -2/5, 1/(q - 2))$

We remark that the existence of these exceptional surgeries was previously known [EM97].

The key fact in our method of analyzing these knots is that they are strongly invertible. Let $\mathcal{T}_{p,q}$ be the tangle obtained by performing the Montesinos trick (see Figure 3.1). We now have the advantage of viewing the surgery space $K_{p,q}(\alpha_r)$ as the branched double cover of S^3




FIGURE 3.1. The pretzel knot $P(-2, 2p + 1, 2q + 1)$, the quotient tangle $\mathcal{T}_{p,q}$, and the pretzel knot $P(-2, 5, -3)$, shown as the boundary of a punctured Klein bottle




FIGURE 3.2. The tangle $\mathcal{T}_{1,q}$, along with fillings $\mathcal{T}_{1,q}(1)$ and $\mathcal{T}_{1,q}(2)$ and the respective Montesinos links that result after isotopy: $K[2/3, -2/5, 1/(q - 2)]$ and $K[1/2, -1/4, 2/(2q - 5)]$

along $\mathcal{T}_{p,q}(r)$. It is easy to verify the two classes of exceptional surgeries in Theorem 1.1 by noticing that $\mathcal{T}_{1,q}(4q + 7)$ and $\mathcal{T}_{1,q}(4q + 6)$ are the Montesinos links $K[2/3, -3/5, 1/(q - 2)]$ and $K[1/2, -1/4, 2/(2q - 5)]$, respectively. [See Figure 3.2.]

The proof that the $K_{p,q}$ admits no other small Seifert fibered surgeries is accomplished by the following two lemmas and the techniques of Subsection 2.7.

Lemma 3.1. If $K_{p,q}(\alpha_r)$ is a small Seifert fibered space for $p \neq 1$, then $|p| \leq 8$ and $|q| \leq 8$.

Lemma 3.2. If $K_{1,q}(\alpha_r)$ is a small Seifert fibered space for $r \notin \{1, 2\}$, then $|q| \leq 8$.

Before we prove these lemmas, we should remark on the possible surgery slopes α_r . We notice that each knot $K_{p,q}$ bounds a punctured Klein bottle at slope α_0 (see Figure 3.1). It follows that $K_{p,q}(\alpha_0)$ is toroidal. By Theorem 2.10, it follows that if $K_{p,q}(\alpha_r)$ is a small Seifert fibered space, then $\Delta(\alpha_r, \alpha_0) \leq 8$.




FIGURE 3.3. The link $\mathcal{T}_{p,q}(0)$, whose branched double cover corresponds to α_0 -surgery on $K_{p,q}$

In many cases, it should be possible to reduce this distance bound to 5, but this is dependent on work in progress by Boyer, Gordon, and Zhang [BGZ12]. However, using Theorem 2.11, we can fairly easily show the following lemma.

Lemma 3.3. If $K_{p,q}(\alpha_r)$ is a small Seifert fibered space, and $|p|, |q| \geq 4$, then $\Delta(\alpha_r, \alpha_0) \leq 6$.

Of course, if r is integral, this means that $|r| \leq 6$, and if we have $r/s \in \mathbb{Q}$, we have that $|4(p+q+1)s - r| \leq 6$.

Proof of Lemma 3.3. We begin by noticing that $K_{p,q}(\alpha_0) = D^2(2, 2) \cup_{T^2} X_{p,q}$ (see Figure 3.3). Under the hypotheses of the lemma, we will show that $X_{p,q}$ is not a Seifert fibered space, so Theorem 2.11 gives us the desired bound. If $p = 1$, then $X_{1,q} = D^2(3, |q-1|)$, and Theorem 2.11 does not apply.

Consider the following fillings on $X_{p,q}$. (See Figure 3.4.)

$$\begin{aligned} X_{p,q}(0) &= L(p+q+1, 1) \\ X_{p,q}(\infty) &= L(4pq - 2p - 2q - 3, 2pq - q - 2) \\ X_{p,q}(-1) &= S^2(1/3, 1/(p-1), q/(q-1)) \end{aligned}$$

If $X_{p,q}$ is a Seifert fibered space, then it has, for its base surface, either D^2 or M^2 (the Möbius band). We will make use of Lemma 2.2 If $X_{p,q}$ is Seifert fibered over the disk with more than two exceptional fibers it cannot have lens space fillings. If $X_{p,q}$ has the form of $D^2(a)$, then it cannot have fillings with three exceptional fibers, so $X_{p,q}(-1)$ must be a lens space. This implies that $p = 2$ or $q = 2$. If $X_{p,q}$ has base surface M^2 , then it can only have lens space fillings or fillings with at least three exceptional fibers, two of which have multiplicity two. Thus, we must have $p, q = 2, 3$. So, assume $X_{p,q} = D^2(a, b)$.

In this case, $X_{p,q}$ has one reducible filling at slope γ and the property that any lens space filling must be at distance one from γ . By considering the three fillings given above, it follows that $\gamma = 0, \infty$, or ± 1 . If $\gamma = -1$, then $X_{p,q}(-1)$ must be reducible, so $p = 1$ or $q = 1$, both of which are not allowed values. If $\gamma = 0$ or if $\gamma = \infty$, then $X_{p,q}(-1)$ must be a lens space, so $p = 2$ or $q = 2$. Finally, if $\gamma = 1$, then the filling $X_{p,q}(-1)$ is at distance two from the reducible filling, so it must have an exceptional fiber of multiplicity 2. It follows that $p = 3$ or $q = 3$. $\square$

We remark that the lemma could be strengthened to say that $X_{p,q}$ is non-Seifert fibered if and only if $p \neq 1$ by showing that $X_{p,q}(1)$ is neither reducible, a lens space, or a small Seifert fibered space with finite fundamental group, as would need to be the case given the different




FIGURE 3.4. The three fillings, 0, ∞ , and 1, on $\mathcal{T}_{p,q}$ used to show that $X_{p,q}$ is not Seifert fibered.

Seifert fibered structures $X_{p,q}$ might have. However, we will not need anything stronger than what we have proved.

Proof of Lemma 3.1. Suppose that $p \neq 1$ and remove a ball around the q -twists of $\mathcal{T}_{p,q}(r)$ to form the tangle $\mathcal{S}_{p,r}$ (see Figure 3.5). Let $N_{p,r}$ denote the branched double cover of $\mathcal{S}_{p,r}$. First, we will show that $N_{p,r}$ is hyperbolic.

We begin by showing some interesting fillings of $N_{p,r}$ (see Figure 3.5).

$$\begin{aligned} N_{p,r}(-1/q) &= K_{p,q}(\alpha_r) \\ N_{p,r}(0) &= D^2(1/2, -1/2) \cup_{T^2} D^2(-1/2, p/(2p-1)) \\ N_{p,r}(\infty) &= T(2, 2p+3)(\alpha_r) = S^2(-1/2, -1/(r+2), -2/(2p+3)) \\ N_{p,r}(-1) &= K[-2p/(6p+1)](\alpha_r) \end{aligned}$$

We remark that $N_{p,r}(\infty)$ and $N_{p,r}(-1)$ correspond to (α_r) -surgery on $K_{p,0}$ and $K_{p,-1}$, respectively. The latter is a 2-bridge knot with no exceptional fillings (if $p \neq 1$), according to the classification by Brittenham and Wu ([BW01]). Thus, $N_{p,r}(-1)$ is hyperbolic if $p \neq 1$.

Now, suppose that $N_{p,r}$ is not hyperbolic, so it must be reducible, ∂ -reducible, Seifert fibered, or toroidal by geometrization. However, $N_{p,r}$ cannot be reducible, since it has two distinct irreducible fillings at slopes 0 and -1 (this follows from the solution to the knot complement problem [GL89]). Similarly, it cannot be Seifert fibered, since it has a hyperbolic filling at slope -1 . It follows that $N_{p,r}$ cannot be ∂ -reducible, since the only irreducible, ∂ -reducible manifold with torus boundary is Seifert fibered, namely, the solid torus.

Finally, suppose that $N_{p,r}$ is toroidal. If any essential torus were non-separating, then all fillings of $N_{p,r}$ would contain an essential non-separating torus, which is false here. Suppose any essential torus is separating, and decompose $N_{p,r}$ along an outermost such torus, F , so that $N_{p,r} = A \cup_F B$ with A atoroidal and $\partial N_{p,r} \subset B$. If we assume, for a contradiction, that $N_{p,r}(-1/q)$ is small Seifert fibered for some q with $|q| > 8$, then we have that F compresses in




FIGURE 3.5. The tangle $\mathcal{S}_{p,r}$, along with two fillings, $\mathcal{S}_{p,r}(0)$ and $\mathcal{S}_{p,r}(\infty)$, and their equivalents after isotopy.

$B(-1)$, $B(\infty)$, and $B(-1/q)$. It follows, from Lemma 2.1 that B is a cable space with cabling slope $\gamma = 0$. But $N_{p,r}(0)$ is neither reducible nor a lens space, so we reach a contradiction. It follows that $N_{p,r}$ is not toroidal, and must be hyperbolic.

Since $N_{p,r}$ is hyperbolic, and $N_{p,r}(\infty)$ is exceptional, it follows that for any exceptional filling $N_{p,r}(-1/q)$, $\Delta(\infty, -1/q) \leq 8$, by Theorem 2.10. It follows that $|q| \leq 8$, as desired. A similar argument shows that $|p| \leq 8$, as well. $\square$

Proof of Lemma 3.2. We will proceed as in the lemma above by analyzing the tangle $\mathcal{S}_r = \mathcal{S}_{1,r}$ formed by removing a ball containing the q -twist region of knot $\mathcal{T}_{1,q}(r)$. We will show that the branched double cover N_r of $\mathcal{S}_r$ is hyperbolic.

Consider the following fillings on N_r (see Figure 3.6). Assume for a contradiction that $q \geq 9$ and $K_{p,q}(\alpha_r) = N_r(-1/q)$ is a small Seifert fibered space.

$$\begin{aligned} N_r(-1/q) &= K_{p,q}(\alpha_r) \\ N_r(0) &= S^2(1/2, -1/2, 1/(2-r)) \\ N_r(\infty) &= S^2(1/2, -2/5, -1/(r+2)) \\ N_r(-1) &= S^2(1/3, -1/4, -1/r) \\ N_r(-1/2) &= S^2(-1/3, 2/5, -1/(r-1)) \\ N_r(1) &= K[-2/7](\alpha_r) \end{aligned}$$

It is clear from this that N_r is irreducible (again, by [GL89]), since it has distinct irreducible fillings, for any value of r . Suppose that N_r is Seifert fibered. Since, for all values of r , N_r has fillings that are Seifert fibered with base surface S^2 , but do not contain a pair of exceptional fibers of multiplicity 2, the base surface of N_r is orientable, i.e., D^2 . A Seifert fibered space with connected boundary with a small Seifert fibered filling must have 2 or 3 exceptional fibers. Furthermore, since no slope is distance one from 0, ∞ , and -1 , N_r has 2 exceptional fibers, i.e., $N_r = D^2(a, b)$.




FIGURE 3.6. Four fillings, 0, ∞ , 1, and $1/2$, of the tangle $\mathcal{S}_r$ that help to prove that N_r is hyperbolic if $r \neq 0, 1, 2$.

By the classification of exceptional surgeries on 2-bridge knots [BW01], $N_r(1)$ is exceptional if and only if $\alpha_r \in \{0, 1, 2, 3, 4\}$. In this case, $p = 1$ and $q = -1$, so $\alpha_r = 4 - r$ and this is the equivalent to $r \in \{4, 3, 2, 1, 0\}$. However, we already know that if $r = 0, 1, 2$, then $N_r(-1/q)$ is exceptional, so we only need to consider $r = 3, 4$.

If $r = 3$, then, by considering $N_3(\infty)$, we see that $a = 5$, and by considering $N_3(-1)$, we see that a cannot be 5. If $r = 4$, then by considering $N_4(-1/2)$, we see that $a = 3$, and by considering $N_4(\infty)$, we see this is impossible. It follows that N_r cannot be Seifert fibered if $r \notin \{0, 1, 2\}$.

It follows that N_r is non-Seifert fibered and, thereby, ∂ -irreducible. If N_r were toroidal, since it has atoroidal fillings at distance two, it must be a cable space, by Lemma 2.1. However, the only cabling slope γ that satisfies $\Delta(\gamma, -1/2) = 1$ and $\Delta(\gamma, \infty) = 1$ is $\gamma = 0$, in which case we must have $N_r(0)$ be reducible or a lens space. So, we must have $r = 3$ and $N_3(0)$ is a lens space. Suppose $N_3 = A \cup_{T^2} B$ with A atoroidal and $\partial N_3 \subset B$.

Then, because $B(\alpha) = S^1 \times D^2$ for $\alpha \in \{-1/q, \infty, -1, -1/2, 1\}$ each of these fillings induces a filling β_α on A . Since $B(0) = (S^1 \times D^2) \# L$ for some lens space L , and since $N_3(0)$ is a lens space, it follows that $A(\beta_0) = S^3$, so A is a knot complement. By construction, A is atoroidal. If A were Seifert fibered, then, by considering $A(\beta_\infty) = N_3(\infty)$ and $A(\beta_{-1}) = N_3(-1)$ just as before, we reach a contradiction.

It follows that N_r must be hyperbolic (for $r \notin \{0, 1, 2\}$). An application of Theorem 2.10 gives us that $\Delta(-1/q, \infty) \leq 8$ if $N_r(-1/q)$ is a small Seifert fibered space, which proves the lemma. □

3.1. Completing the proof of Theorem 1.1. The work above leaves us with a finite list of knots and links $L_{p,q,r} = \mathcal{T}_{p,q}(r)$ whose branched double covers might be Seifert fibered. We must consider non-integral r if and only if p and q are both positive. By Proposition 2.6, we must show that each of these links is not a Montesinos link or a Seifert link.

Method 1 (see Subsection 2.7) can be used to show that none of the $L_{p,q,r}$ are Montesinos knots or links, though it should be noted that the Kauffman polynomial must be employed in

a handful of cases, including distinguishing $L_{1,4,-1}$ from $K[1/3, 2/5, -2/5]$ and for some non-integral values of r , and that the HOMFLYPT polynomial must be employed to distinguish $L_{1,6,-2}$ from $K[-1/4, 1/6, 2/7]$. In other words, these pairs are not distinguished by their Alexander polynomial or Khovanov homology.

Now, consider when $L_{p,q,r}$ is a link. Then it is the union of the unknot and the 2-bridge knot $K[n/m]$, where $n = 2pq - p - 2$ and $m = 4pq - 2p - q - 3$. $K[n/m]$ is a torus knot only if $p = 1$ and $q = 3$ or 4 . In the latter case, $L_{1,4,r}$ is the union of a trefoil and an unknot. If this link is to be a Seifert link, the unknotted component must lie as the core of the torus upon which the trefoil sits. However, the link just described can be distinguished from $L_{1,4,r}$ for all values of r using the Jones polynomial. Concerning $L_{1,3,r}$, exceptional surgeries on the knot $P(-2, 3, 7)$ are previously well-understood [EM97].

It only remains to show that $L_{p,q,r}$ is never a torus knot. If $p \neq 1$, this is accomplished by applying Criterion 2.17. For $p = 1$, Criterion 2.18 suffices.

4. THE CASE OF $(3, 3, |q_3|)$

We now turn our attention to pretzel knots $P(q_1, q_2, q_3)$ such that $|q_1| = |q_2| = 3$. The case of $P(3, 3, q_3)$, where $q_3 > 0$ was handled by Ichihara and Jong in [IJ11]. We break up the remaining cases as follows.

- (1) $P(3, \pm 3, -2m)$ with $m \geq 1$
- (2) $P(3, 3, 2m + 1)$ with $m \leq -2$
- (3) $P(3, -3, 2m + 1)$ with $m \leq -3$ or $2 \leq m$
- (4) $P(3, 3, 2m, -1)$ with $2 \leq m$

In Cases (2) and (3), it is only necessary to consider integral surgery slopes by Theorem 1.5. Our main result is:

Theorem 1.2. Hyperbolic pretzel knots of the form $P(3, 3, m)$ or $P(3, 3, m, -1)$ admit no small Seifert fibered surgeries. Hyperbolic pretzel knots of the form $P(3, -3, m)$ admit small Seifert fibered surgeries precisely in the following cases:

- $P(3, -3, -2)(-1) = S^2(2/3, -1/4, -2/5)$
- $P(3, -3, -3)(-1) = S^2(-1/2, 1/5, 2/7)$
- $P(3, -3, -4)(-1) = S^2(1/2, -1/5, -2/7)$
- $P(3, -3, -5)(-1) = S^2(1/3, 1/4, -3/5)$
- $P(3, -3, -6)(-1) = S^2(1/2, -1/3, -2/13)$

4.1. Case (1). Let $K_m^\pm = P(3, \pm 3, -2m)$. To avoid the redundancy of mirrors, we can restrict to $m > 0$. Recall that these knots can only admit non-integral small Seifert fibered surgeries in the case of K_m^+ . Our first result is half of Theorem 1.2.

Proposition 4.1. Let $K_m^\pm = P(3, \pm 3, -2m)$, with $m > 0$, be hyperbolic. Then, $K_m^\pm$ admits no small Seifert fibered surgeries, except in the following three instances.

- $P(3, -3, -2)(-1) = S^2(2/3, -1/4, -2/5)$
- $P(3, -3, -4)(-1) = S^2(1/2, -1/5, -2/7)$
- $P(3, -3, -6)(-1) = S^2(1/2, -1/3, -2/13)$

As in Section 3, we will proceed in this case by first limiting the possible surgery slopes, then limiting the size of m , then using techniques from Subsection 2.7 to check that small values of m and slopes satisfying the relevant bound do not produce small Seifert fibered spaces (except for the three noted cases). Let $K_m^\pm = P(3, \pm 3, -2m)$ with $m > 0$. We begin by




FIGURE 4.1. (a) The knot $P(3, -3, -2m)$ (shown here with $m = 3$) bound by punctured Klein bottles. (b) The tangle $\mathcal{T}_m^+$.




FIGURE 4.2. The tangle $\mathcal{S}_r^- r$ and the link $L_r(m)$

observing that $K_m^\pm$ bounds a punctured Klein bottle. Let $\alpha_r^+ = 12 - r$ and $\alpha_r^- = -r$ (again, these are chosen so that $K_m^\pm(\alpha_r)$ corresponds to r -filling on the corresponding tangle). Then this Klein bottle has boundary slope $\alpha_0^\pm$ (see Figure 4.1(a) for the case of K_m^-). Since surgery along this slope produces a toroidal manifold (as in the previous section), any exceptional surgery slope for $K_m^\pm$ must be close to $\alpha_0^\pm$. In particular, $\Delta(\alpha_r^\pm, \alpha_0^\pm) \leq 8$.


Lemma 4.2. Suppose $K_m^\pm(\alpha_r)$ is a small Seifert fibered space. Then, $m \leq 8$.

Proof. Let $L_r^\pm(-m) = \mathcal{T}_m^\pm(r)$ and form the tangle $\mathcal{S}_r^\pm$ by removing a 3-ball containing the m -twist box of $L_r^\pm(-m)$ (see Figure 4.2). Let $M_r^\pm = \widetilde{\mathcal{S}_r^\pm}$. Of course, $\mathcal{S}_r^\pm(-1/m) = L_r^\pm(-m)$, and $K(\alpha_r) = M_r^\pm(1/m)$. Assume, for a contradiction, that $M_r^\pm(1/m)$ is a small Seifert fibered space for some $m \geq 9$.

Consider the following fillings of $M_r^\pm$, which we can easily visualize and verify by looking at the corresponding rational tangle fillings of $\mathcal{S}_r^\pm$ (see Figure 4.3).

$$\begin{aligned} M_r^\pm(1/m) &= \text{small Seifert fibered space (by assumption)} \\ M_r^-(0) &= (S^1 \times S^2) \# L(r, 1) \\ M_r^+(0) &= D^2(2, 2) \cup_{T^2} D^2(2, r) \\ M_r^\pm(\infty) &= D^2(2, 3) \cup_{T^2} D^2(2, 3) \end{aligned}$$

As was argued in Section 3, $M_r^\pm$ is irreducible (it has distinct irreducible fillings), non-Seifert fibered (it has a non-Seifert fibered, non-reducible filling), and ∂ -irreducible (it is irreducible and not $S^1 \times D^2$). Assume that M_r^- is toroidal, so $M_r^- = A \cup_F B$ with A atoroidal and $\partial M_r^- \subset B$.

FIGURE 4.3. Two interesting fillings of $\mathcal{T}_r$

Suppose that F compresses in $M_r^-(\infty)$. Then, since $\Delta(1/m, \infty) = |m| \geq 2$, B is a cable space with cabling slope γ satisfying $\Delta(\gamma, \infty) = \Delta(\gamma, 1/m) = 1$. It follows that $\gamma = a \in \mathbb{Z}$, and $|ma - 1| = 1$. Since $|m| \geq 9$, a must be zero, so $\gamma = 0$. It follows that $B(0) = (S^1 \times D^2) \# L$, where L is a lens space, and $B(\infty) = B(1/m) = S^1 \times D^2$. Let β_0, β_∞ , and $\beta_{1/m}$, be the slopes of the induced slopes of the meridian of B after the above fillings are performed, so $M_r^-(\alpha) = A(\beta_\alpha)$ for $\alpha = \infty$ or $1/m$, and $M_r^-(0) = A(\beta_0) \# L$. Since L has finite fundamental group, $L = L(r, 1)$ and $A(\beta_0) = S^1 \times S^2$.

Now, since $\Delta(\beta_\infty, \beta_0) \geq 4$ (by Lemma 2.1), A cannot be hyperbolic, because $\Delta(S^2, T^2) = 3$. Since A was assumed to be atoroidal, it follows that A must be Seifert fibered. But $A(\beta_\infty)$ is irreducible and not Seifert fibered, so this cannot be. This contradiction means that F cannot compress in $M_r^-(\infty)$.

So, assume F remains incompressible in $M_r^-(\infty)$. If $|r| = 1$, then $M_1^-(0) = S^1 \times S^2$. But since F is the unique incompressible torus in a non-Seifert fibered graph manifold, $A = D^2(2, 3)$. In particular, $M_1^-(0) = A(\beta_0) = S^1 \times S^2$ is not a possible filling of a trefoil complement.

If $|r| > 1$, then since F compresses in fillings at slopes $1/m$ and 0 , which are at distance one, by Lemma 2.1, B is either a cable space or the exterior of a braid in a solid torus. Since $M_r^-(0) = (S^1 \times S^2) \# L(r, 1)$, either $B(0)$ or $A(\beta_0)$ is $S^1 \times S^2$. However, this is not possible for such spaces B , nor is it possible for $A = D^2(2, 3)$.

Thus, M_r^- is not toroidal, and must be hyperbolic. So, by Theorem 2.11, $\Delta(1/m, \infty) = |m| \leq 8$, a contradiction that yields the desired result. The reasoning is very similar to show that M_r^+ must be hyperbolic as well, noting that $M_r^+(0)$ is hyperbolic, regardless of r . $\square$

4.2. Completing the proof of Proposition 4.1. By our work above, we can conclude that if $P(3, \pm 3, -2m)(\alpha_r^\pm)$ with $m > 0$ admits a small Seifert fibered surgery, then $|r|, |m| \leq 8$. Let $L_{m,r}^\pm = \mathcal{T}_m^\pm(r)$. We assume that r is integral for $L_{m,r}^-$.

First, consider the links $L_{m,r}^\pm$, which are the union of an unknotted component with a component $J_m^\pm = K[2/3, \pm 2/3, -1/2m]$. Because $J_m^\pm$ is not a torus knot or a 2-bridge knot




FIGURE 4.4. The left figure above shows the pretzel knot $P(3, 3, 2m + 1)$ as the boundary of a punctured torus, while the right one exhibits the strong inversion of $P(3, 3, 2m + 1)$ given by rotation. Here, $m = -3$.

for any m , by Criteria 2.15 and 2.19, we can conclude that $L_{m,r}^\pm$ is never a 2-component Seifert link or Montesinos link.

When $L_{m,r}^\pm$ is a knot, we see that $2g(L_{m,r}^\pm) \neq 2(L_{m,r}^\pm)$, so, by Criterion 2.17, $L_{m,r}^\pm$ is never a torus knot. To see that $L_{m,r}^\pm$ is never a Montesinos knot, we implement Method 1 (see Subsection 2.7), accounting for r non-integral when necessary.

4.3. Case (2). Next, we will consider hyperbolic pretzel knots of the form $K_m = P(3, 3, 2m + 1)$. Here, K_m is hyperbolic if $m \neq -1$ or 0 , and if m is positive, then Ichihara-Jong have shown that K_m admits no Seifert fibered surgeries [IJ11]. The case when $m = -2$ will be covered in Subsection 4.5, so assume $m \leq -3$. In this section, we prove the following.

Proposition 4.3. A hyperbolic pretzel knot of the form $P(3, 3, 2m + 1)$ with $m \leq -3$ admits no small Seifert fibered surgeries.

As before, we will first restrict the possible values of m for which K_m might admit a Seifert fibered surgery, then rule the remaining cases out by computer. First we note that K_m has genus one, so by Theorem 2.12, $|r| \leq 3$ (see Figure 4.4). Since the three pretzel parameters are all odd, K_m cannot admit non-integral Seifert fibered surgeries by Theorem 1.5. Let $\alpha_r = -r$, so that $K_m(\alpha_r) = K_m(-r)$ will correspond with $\mathcal{T}_m(r)$ (see below), as before.

Lemma 4.4. If $m \leq -10$, then $K_m(r)$ is not a small Seifert fibered space.

Proof. In a slight variation of the preceding cases, these knots possess a strong inversion that is a half-rotation of the plane. Let $\mathcal{T}_m(r)$ be the resulting quotient link, as before (see Figure 4.4). Again, we form the tangle $\mathcal{S}_r$ by removing a ball containing the $(m + 1)$ -twist region. If we denote by N_r the branched double cover of S^3 along $\mathcal{S}_r$, and assume for a contradiction that $K_m(r)$ is a small Seifert fibered space for some $m \leq -10$ and some r , then we have the following fillings of N_r (see Figure 4.6). Note that $N_r(\infty) = P(3, 3, -1)(r)$, so it is simply r -surgery on the left-handed trefoil.

$$\begin{aligned} N_r(-1/(m + 1)) &= \text{small Seifert fibered space (by assumption)} \\ N_r(0) &= \text{non-Seifert fibered toroidal space} \\ N_r(\infty) &= S^2(-1/2, 1/3, -1/(r + 6)) \end{aligned}$$

Since N_r has distinct irreducible fillings and a non-Seifert fibered irreducible filling, N_r is irreducible, non-Seifert fibered, and ∂ -irreducible. If N_r is toroidal, then since it has atoroidal fillings at distance $\Delta(-1/(m + 1), \infty) = |m + 1| > 2$, it has as a subspace a cable space with




FIGURE 4.5. The figures above illustrate how to obtain the quotient tangle $\mathcal{T}_m$ by applying the Montesinos trick to the strong inversion of $P(3, 3, 2m + 1)$ given by rotating the knot π radians through its center. Here, $m = -3$.




FIGURE 4.6. The first figures above, from left to right, are: (a) the link $\mathcal{T}_m(-\alpha_r)$, (b) the tangle $\mathcal{S}_r$, the filling $\mathcal{S}_r(\infty)$, which is isotopic to $K[-1/2, 1/3, 1/(6 - r)]$, and (d) the fillings $\mathcal{S}_r(0)$, whose link complement contains an essential torus.

cabling slope $\gamma = 0$. This means that $N_r(0)$ is either reducible or a lens space, which is a contradiction. It follows that N_r is hyperbolic and that $\Delta(-1/(m + 1), \infty) = |m + 1| \leq 8$, a contradiction that completes the proof. $\square$

4.4. Completing the proof of Proposition 4.3. Let $L_{m,r} = \mathcal{T}_m(r)$ denote the quotient link described above, and note that it is only necessary to consider $r \in \mathbb{Z}$ here. If $L_{m,r}$ is a link, then we see that it is the union of a trefoil with the knot $J_m = K[1/2, 1/3, (m - 1)/(2m - 3)]$.




FIGURE 4.7. (a) The knot $P(3, -3, 2m+1)$ (here, $m = -3$), shown as the boundary of a punctured torus, (b) along with the axis of the cycle f of period 2 of the knot, and (c) the quotient knot K_f , here, the unknot, along with the image of $\text{Fix}(f)$ in the quotient.

Since $m \leq -3$, by assumption, J_m is never a torus knot or a 2-bridge knot. It follows that $L_{m,r}$ is never a Seifert link or a Montesinos link, by Criteria 2.19 and 2.15, respectively.

When $L_{m,r}$ is a knot, we see that $|\text{Kh}(L_{m,r})| = 6$ and $s(L_{m,r}) < \text{br}(\Delta_{K_{m,r}}(t))$, so $L_{m,r}$ cannot be a Montesinos knot or a torus knot by Criteria 2.14 and 2.17, respectively.

4.5. Case (3). We now consider hyperbolic pretzel knots $K_m = P(3, -3, 2m+1)$. We will allow m to be positive or negative, which will allow us to consider only positive surgery slopes (which must be integral if they are to be exceptional by Theorem 1.5). These are genus one knots, so $K_m(r)$ can only be exceptional if $|r| \leq 3$ by Theorem 2.12. Thus, we assume $r = 1, 2$, or 3 .

These knots are the first that we have encountered with no strong inversion, so we cannot make use of the Montesinos trick. However, K_m does have cyclic period 2, so we will study the space $K_m(r)$ by studying the link $(L_m)_f$, which is the image of $\text{Fix}(f)/\langle f \rangle$ in $(K_m)_f(r/2)$ (recall this set-up from Section 2.5). See Figure 4.7.

Proposition 4.5. The hyperbolic pretzel knot $P(3, -3, 2m+1)$ admits a small Seifert fibered surgery precisely in the following instances.

- $P(3, -3, 3)(1) = S^2(1/2, -1/5, -2/7)$
- $P(3, -3, 5)(1) = S^2(-1/3, -1/4, 3/5)$

Note that these exceptional surgeries are presented as the mirrors of those presented in Theorem 1.2. Again, our first task is to restrict the possible values of m for which K_m might admit a Seifert fibered surgery, then rule the remaining cases out by computer. We will handle the three cases $r = 1, 2$, and 3 separately below.

Lemma 4.6. The space $P(3, -3, 2m+1)(1)$ is not a small Seifert fibered space for $|m| \geq 9$.

Proof. Assume that $P(3, -3, 2m+1)(1)$ is a small Seifert fibered space with $|m| \geq 9$. As we have seen $K_m(1)$ is the branched double cover of S^3 along the knot $(L_m)_f$. We form the tangle $\mathcal{S}$ by removing the m -twist box of $(L_m)_f$. Let $Z = \tilde{\mathcal{S}}$ (see Figure 4.8). Then, we have the following fillings:

$$\begin{aligned}
 Z(-1/m) &= \text{small Seifert fibered (by assumption)} \\
 Z(0) &= S^2 \times S^1 \\
 Z(-1/3) &= D^2(2, 3) \cup_{T^2} D^2(2, 3) \\
 Z(-1/2) &= S^2(3, 5, 7) \\
 Z(-1) &= S^2(2, 5, 7) \\
 Z(\infty) &= S^2(2, 3, 11)
 \end{aligned}$$




FIGURE 4.8. (a) The link $K_f \cup \text{Fix}(f)$ in the quotient, (b) the knot L_f resulting from $(1/2)$ -surgery on the unknotted component, and (c) the tangle $\mathcal{S}$ formed by removing the m -twist area of the knot. Here, $m = -3$.




FIGURE 4.9. Above, we see five interesting fillings of $\mathcal{S}$: $\mathcal{S}(1)$, $\mathcal{S}(1/2)$, $\mathcal{S}(1/3)$, $\mathcal{S}(0)$, and $\mathcal{S}(\infty)$.

It is worth noting that the last four fillings on the list correspond to exceptional fillings of hyperbolic pretzel knots. $P(3, -3, 7)(1) = Z(-1/3)$, $P(3, -3, 5)(1) = Z(-1/2)$, $P(3, -3, 3)(1) = Z(-1)$, and $P(3, -3, 1)(1) = Z(\infty)$. The lattermost is surgery on the rational knot $\mathcal{K}[-2/9]$. See Figure 4.9.

Because Z has distinct irreducible fillings as well as an irreducible non-Seifert fibered filling, it is impossible for Z to be Seifert fibered, reducible, or ∂ -reducible. Assume that Z is toroidal, so $Z = A \cup_F B$ with A atoroidal and $\partial Z \subset B$. Then, since F compresses in $Z(-1/2)$, $Z(-1)$, and $Z(\infty)$, and $\Delta(1/2, \infty) \geq 2$, B must be a cable space. The cabling slope γ is restricted to be distance one from ∞ and $-1/2$, so $\gamma = 0$ or -1 . Since $Z(-1)$ is not a lens space nor reducible, we cannot have $\gamma = -1$. If $\gamma = 0$, then $B(0) = (S^1 \times D^2) \# (S^1 \times S^2)$, which is not a possible result of filling on a cable space.

It follows that Z is hyperbolic, so $\Delta(-1/m, \infty) \leq 8$, so $|m| \leq 8$, which gives the desired contradiction. □

Proposition 4.7. Let $P(3, -3, 2m+1)$ be a hyperbolic pretzel knot. Then, $P(3, -3, 2m+1)(2)$ is never a small Seifert fibered space.




FIGURE 4.10. (a) The link $K_f \cup \text{Fix}(f)$ in the quotient, (b) the link L_f resulting from 1-surgery on the unknotted component (note that the core of the surgery torus is a component of the resulting link), and (c) the tangle formed by removing the m -twist area of the knotted component J_m of the resulting link. Here, $m = -3$.

Proof. We can proceed as above, by analyzing the result of 1-surgery on the quotient knot $(K_m)_f$. This gives a two-component link in S^3 , $(L_m)_f$, such that the double cover of S^3 branched along $(L_m)_f$ is the surgery space $P(3, -3, 2m+1)(2)$. Thus, to show this surgery space is not a small Seifert fibered space, it suffices to show that $(L_m)_f$ is neither a Montesinos link of length three with two components nor a Seifert link with two components.

First, we note that $(L_m)_f$ is the union of the unknot with the knot $J_m = K[1/3, -1/3, 1/(m-1)]$ (see the right half of Figure 4.10). Since J_m is never torus knot, by Criterion 2.19, $(L_m)_f$ is never a Seifert link. If $m = 0$ or $m = 2$, then $J_0 = K[-2/9]$ and $J_2 = K[2/9]$, respectively; otherwise, J_m is not a 2-bridge knot, so $(L_m)_f$ is not a Montesinos link, by Criterion 2.19.

If $m = 0$, then we are considering 2-surgery on $P(3, -3, 1)$, which is $K[2/9](2)$. By the classification of Brittenham and Wu [BW01], this space is Seifert fibered. If $m = 2$, then we are considering the space $P(3, -3, 5)(2)$, and if $(L_m)_f$ is a Montesinos link, then it has the form $K[x, y, z]$, where $z = 2/9$ or $4/9$ and x and y have even denominator.

Now, since $(L_2)_f$ has a diagram with 12 crossings, and since the z -tangle would contribute 6 crossings, if $(L_2)_f$ is to be a two component length-3 Montesinos link, then the x - and y -tangles must contribute at most 6 crossings. Without loss of generality, we can assume $x = \pm 1/2$ and $y = \pm 1/2$ or $\pm 1/3$ or $\pm 3/4$. However, an easy check shows that the determinant of such Montesinos links cannot be 2.

□

Finally, we consider the case of 3-surgery on hyperbolic pretzel knots K_m of the type $P(3, -3, 2m+1)$. These knots have period 2, so we can analyze the surgery space $K_m(3)$ as the double branched cover of $(K_m)_f(3/2)$, where $(K_m)_f$ is the factor knot K_m/f for the self diffeomorphism $f : S^3 \rightarrow S^3$ of order two that preserves K_m . In this case, $(K_m)_f$ is the unknot, so $(K_m)_f = -L(3, 2)$. Let L_m denote the image of $\text{Fix}(f)/f$ in the surgery space $(K_m)_f(3/2)$, i.e., L_m is the branching set for the double covering.

Proposition 4.8. Let $P(3, -3, 2m+1)$ be a hyperbolic pretzel knot. Then $P(3, -3, 2m+1)(3)$ is never a small Seifert fibered space.

Proof. By the analysis of [MM02], the link L_m is actually a 2-bridge link, contained in the solid torus that, together with the surgery solid torus, comprises half of the genus-1 Heegaard splitting of the lens space $-L(3, 2)$ (see Figure 4.11). Thus, if we pass to a 3-fold cover, the




FIGURE 4.11. Above, on the left we have the link $(K_m)_f \cup \text{Fix}(f)/f$ in the quotient, and, on the right, we have the result of $(3/2)$ -surgery on the unknotted component: the link L_m contained in a solid torus (simply view the knot as lying in the solid torus that comprises the exterior of unknot).




FIGURE 4.12. The link $\widetilde{L}_m$ in S^3 , which is the triple cover of L_m in $-L(3, 2)$.

lift, $\widetilde{L}_m$, of L_m will be a length-3 Montesinos link, contained in one half of the standard genus-1 Heegaard splitting of S^3 . We now describe how to see this.

The two solid tori that comprise the splitting of $-L(3, 2)$ are attached via a map which sends the meridian of one to a $(-3/2)$ -curve on the boundary of the other. Passing to the 3-fold cover changes the image of the attaching map to a $(-1/2)$ -curve, which gives S^3 . This lift simply triplicates the knotted part of L_m . However, if we want to think about this lift as a knot in the standard 3-sphere, we must apply a self-diffeomorphism of S^3 to get the standard Heegaard splitting of S^3 (i.e., where the meridian of one torus is glued along a $(1/0)$ -curve). This final step introduces two full negative twists of the strands of L_m . The result is a link $\widetilde{L}_m$ in S^3 . See Figure 4.12.

Since $P(3, -3, 2m + 1)(3)$ has quotient $-L(3, 2)$, it has the form $S^2(3, 3, c)$, where the exceptional fiber of multiplicity c corresponds to the branching locus, (L_m) . It follows that the lift $\widetilde{L}_m$ should be a Montesinos link of type $S^2(c, c, c)$. From this, it follows that c^2 divides the determinant of $\widetilde{L}_m$. We can calculate the determinant of this lift to be 49, independent of m , so it follows that $c = 7$, but then $P(3, -3, 2m + 1)(3)$ is $S^2(7, 7, 7)$, which cannot have Euler number 3 (since 7 divides the Euler number), as is required of a space resulting from 3-surgery. $\square$

4.6. Completing the proof of Proposition 4.5. It remains to show that the knots $L_m = \mathcal{T}_m(1)$ are neither Montesinos knots, nor Seifert knots, for $m \in [-8, 8] \setminus \{-1, 0, 1, 2\}$ (notice, when $m = 1, 2$ we *do* get small Seifert fibered surgeries, and when $m = -1, 0$, we have $P(3, -3, \pm 1)$, which are not true pretzel knots).




FIGURE 4.13. The Montesinos knot $P(3, 3, 2m, -1)$ in (a) standard form and (b) pillowcase form, and (c) the factor knot resulting from rotation about the axis perpendicular to the page.




FIGURE 4.14. The pretzel knot $P(3, 3, 2m, -1)$, shown with the axis of its strong inversion, and the resulting tangle $\mathcal{T}_m$. Here $m = 3$.

In fact, for these knots we have that $s(L_m) \neq 2g(L_m)$, so they cannot be torus knots by Criterion 2.17. Furthermore, we can apply Method 1 to show that L_m is never a Montesinos knot.

4.7. Case (4). Finally, consider the case when K_m is a hyperbolic pretzel knot of the form $P(3, 3, 2m, -1)$ with $m > 1$. We note that such knots are often considered to be non-pretzel Montesinos knots. If $m = 1$, then $K_1 = P(-2, 3, 3)$, and is not hyperbolic. We see that K_m has a cyclic of period 2 (see Figure 4.13) with factor knot $K_f = T_{2,3}$ as well as a strong inversion. Since $K_f(r/2)$ must be a lens space surgery on the trefoil, it follows that $\Delta(r/2, 6) = 1$. In this case we consider the possibility of non-integral exceptional surgeries. Here, we note that the quotient tangle $\mathcal{T}_m$ has $4m + 10$ twists, so $L_{m,r} = \mathcal{T}_m(-r)$ has $4m + 10 - r$. See Figure 4.15.

Proposition 4.9. The hyperbolic pretzel knot $P(3, 3, 2m, -1)$ admits no small Seifert fibered surgeries.

Proof. We begin by noting that we can show that $m \leq 8$ just as we did when dealing with K_m^+ , earlier in this section (recall, Figure 4.3), so we will omit the details. Consider the quotient links $L_{m,r}$ obtained via the Montesinos trick. When r is even, they consist of an unknotted component, together with a component J_m , which is the knot $K[-1/3, -1/3, 1/m]$ (see Figure 4.15).

Since J_m is never a torus knot or a 2-bridge knot (for $m > 1$), $L_{m,r}$ is never a Seifert link or a Montesinos link, by Criteria 2.19 and 2.15, respectively.

When $L_{m,r}$ is a knot, we see that $|\text{Kh}(L_{m,r})| \geq 4$ and $s(L_{m,r}) < \text{br}(\Delta_{L_{m,r}}(t))$, so, by Criteria 2.14 and 2.17, $L_{m,r}$ is never a Montesinos knot or a torus knot.

□




FIGURE 4.15. The link $L_{m,r}$ and the component J_m . Here $m = 3$ and $r = 12$ (hence, $4m + 10 - r = 10$ twists).

5. THE CASE OF $(3, 4, 5)$

We next turn our attention to the pretzel knots $P(3, \pm 4, \pm 5)$. We will follow the same program in which we make use of the strong inversion and analyze the quotient link along with its double branched cover to achieve the following result. Because this is not an infinite family of knots, we do not need to argue to restrict any parameters as we have above. Since these knots bound punctured Klein bottles at slope α_0 , we only need to consider fillings $\alpha_r = \alpha_0 - r$ at distance at most 8. Our only task here is to show that the quotient links are not Seifert links nor Montesinos links. In the case of $P(3, -4, 5)$, we must consider non-integral surgeries.

Theorem 1.3. The hyperbolic pretzel knots $P(3, \pm 4, \pm 5)$ admit no small Seifert fibered surgeries.

Proof. Let $L_{\pm, \pm, r}$ be the quotient link resulting from the Montesinos trick. When r is even, $L_{\pm, \pm, r} = U \cup J$, where $J = K[2/3, \pm 1/2, \pm 2/5]$, which is never the unknot, a two-bridge knot, or a torus knot. Thus, by Criteria 2.15 and 2.19, $L_{\pm, \pm, r}$ is never a Montesinos link with two components or a Seifert link with two components.

When $L_{\pm, \pm, r}$ is a knot, $2g(L_{\pm, \pm, r}) \neq s(L_{\pm, \pm, r})$, so $L_{\pm, \pm, r}$ is not a torus knot, by Criterion 2.17, and we can use Method 1 to show that $L_{\pm, \pm, r}$ is never a Montesinos knot. □

6. NON-PRETZEL MONTESINOS KNOTS

In this section we discuss small Seifert fibered surgery on non-pretzel Montesinos knots. By Wu [Wu10, Wu11c], we need only consider a handful of cases: $K[1/3, -2/3, 2/5]$, $K[-1/2, 1/3, 2/(2a+1)]$ for $a \in \{3, 4, 5, 6, \dots\}$, and $K[-1/2, 2/5, 1/(2q+1)]$ for $q \geq 1$. We prove the following theorem.

Theorem 1.4. Suppose that K is a non-pretzel Montesinos knot and $K(\alpha)$ is a small Seifert fibered space. Then either $K = K[-1/2, 2/5, 1/(2q+1)]$ for $q \geq 5$, or K is on the following list and has the described surgeries.

- $K[1/3, -2/3, 2/5](-5) = S^2(1/4, 2/5, -3/5)$
- $K[-1/2, 1/3, 2/7](-1) = S^2(1/3, 1/4, -4/7)$
- $K[-1/2, 1/3, 2/7](0) = S^2(1/2, 3/10, -4/5)$
- $K[-1/2, 1/3, 2/7](1) = S^2(1/2, 1/3, -16/19)$
- $K[-1/2, 1/3, 2/9](2) = S^2(1/2, -1/3, -3/20)$
- $K[-1/2, 1/3, 2/9](3) = S^2(1/2, -1/5, -3/11)$

- $K[-1/2, 1/3, 2/9](4) = S^2(-1/4, 2/3, -3/8)$
- $K[-1/2, 1/3, 2/11](-2) = S^2(-2/3, 2/5, 2/7)$
- $K[-1/2, 1/3, 2/11](-1) = S^2(-1/2, -2/7, 2/9)$
- $K[-1/2, 1/3, 2/5](3) = S^2(1/2, -1/3, -2/15)$
- $K[-1/2, 1/3, 2/5](4) = S^2(1/2, -1/6, -2/7)$
- $K[-1/2, 1/3, 2/5](5) = S^2(-1/3, -1/5, 3/5)$
- $K[-1/2, 1/5, 2/5](7) = S^2(1/2, -1/5, -2/9)$
- $K[-1/2, 1/5, 2/5](8) = S^2(-1/4, 3/4, -2/5)$
- $K[-1/2, 1/7, 2/5](11) = S^2(-1/3, 3/4, -2/7)$

Question 1.8. Do the Montesinos knots $K[-1/2, 2/5, 1/(2q+1)]$ with $q \geq 5$ admit small Seifert fibered surgeries?

With the exception of the case noted in the question above, we now prove the list give in Theorem 1.4 is complete. In proving the theorem, we will consider the three types of Montesinos knots involved separately. Note that throughout, we assume $r \in \mathbb{Z}$.

6.1. The case of $K[1/3, -2/3, 2/5]$. First, consider the case when $K = K[1/3, -2/3, 2/5]$. By Wu [Wu11a], $K(-4)$ and $K(-6)$ are toroidal, so it suffices to consider $K(r)$ for $-12 \leq r \leq 2$, by Theorem 2.10. Define L_r , as we have done before (see Figure 6.1 (left side)). To show that L_r is not a Montesinos link or knot, we apply Method 1. In the case of even r , we note that $L_r = U \cup T(2, 3)$, so if L_r is a length-3 Montesinos link with two components, then one tangle is either a $(1/3)$ -tangle or a $(2/3)$ -tangle. When such a check is performed, precisely one match is found: $K[1/3, -2/3, 2/5](-5)$ is a Seifert fibered space, as shown in Theorem 1.4.

To see that L_r is never a Seifert link, we simply note that for odd r , L_r cannot be a torus knot, by Criterion 2.17, since $|s(L_r)| < \text{br}(\Delta_{L_r}(q))$. If r is even, we note that $L_r = U \cup T(2, 3)$. *A priori*, L_r may be a trefoil union one of its core curves; however, this link has crossing number 7, and the crossing number of L_r is at least $\text{br}(V_{L_r}) = 10$. Thus, L_r is never a two component Seifert link.

6.2. The case of $K[-1/2, 1/3, 2/(2a+1)]$. In the case where $K_a = K[-1/2, 1/3, 2/(2a+1)]$ for $a \in \{3, 4, 5, 6\}$, we note that by Wu [Wu11a] we have the following toroidal fillings: $K_3(-2)$, $K_4(5)$, $K_5(2)$, $K_5(-3)$, and $K_6(2)$, so we consider surgery slopes r with distance at most 8 from the corresponding toroidal filling.

We proceed as above, considering links and knots $L_{a,r}$ (see Figure 6.1 (right side)), to show that $L_{a,r}$ is never a Montesinos knot or link, noting in this case that $L_{a,r} = U \cup J$, where J is $T(2, 5)$ if $a = 3$, $T(2, 3)$ if $a = 4$, and the unknot if $a = 5$ or 6 . By applying Method 1, we find that the Montesinos links are precisely those stated in Theorem 1.4.

If r is even, and $L_{a,r}$ is a Seifert link, we must have that $L_{3,r}$ is the union of $T(2, 5)$, together with a core curve, $L_{4,r}$ is the union of $T(2, 3)$, together with a core curve, and $L_{5,r}$ and $L_{6,r}$ have the form $T(2, 2n)$ for some n . However, $L_{5,r}$ and $L_{6,r}$ cannot have this form, since they are not alternating, a fact deduced by noticing that $|\text{Kh}(L_{a,r})| \geq 3$. As above, $L_{4,r}$ cannot have the said form because it must have at least 9 crossings. Finally, we see that $L_{3,r}$ has linking number ± 1 , while $T(2, 5)$ has linking number ± 5 or ± 2 with its cores. Thus, $L_{a,r}$ cannot be a two component Seifert link.

If r is odd, we can obstruct most of the $L_{a,r}$ from being torus knots by using Criterion 2.17. However, $L_{3,1}$ and $L_{4,1}$ have equal Rasmussen invariant and breadth of Alexander polynomial.




FIGURE 6.1. On the left, we have the Montesinos knot $K[1/3, 1/3 - 3/5]$, along with the axis of one of its strong inversions, and the quotient link L_r . On the right, we have the Montesinos knot $K[-1/2, 1/3, 2/9]$, along with the axis of its strong inversion, and the quotient link $L_{a,r}$ with $a = 4$. In the case of $K[-1/2, 2/5, 1/(2a + 1)]$, we have a picture similar to that on the right.

The former, in fact, corresponds to a Seifert fibered space, so consider $L_{4,1}$. Since this knot has determinant 1, if it is to be a torus knot of the form $T(a, b)$, both a and b are odd. Furthermore, since it has Rasmussen invariant equal to 10, we have $10 = (a - 1)(b - 1)$, which is impossible if a and b are both odd. Thus, $L_{a,r}$ is never a torus knot.

6.3. The case of $K[-1/2, 2/5, 1/(2q + 1)]$. Finally, we are left with the case when $K_q = K[-1/2, 2/5, 1/(2q + 1)]$ and $q \geq 1$. By Wu [Wu11a], we have the following toroidal fillings: $K_1(6), K_2(9), K_3(12), K_4(15)$ so for $q \leq 4$, we can bound the surgery slope r , and complete the classification. For larger q , we cannot. Furthermore, we obtain no bound on q , as we have done above.

If we again consider $L_{q,r}$, we see that for $q \leq 4$ and odd r , many $L_{q,r}$ have $s(L_{q,r}) < \text{br}(\Delta_{L_{q,r}}(t))$ or $\det(L_{q,r}) > s(L_{q,r}) + 1$, so they cannot be torus knots, by Criteria 2.17 and 2.18, respectively. However, $\det(L_{1,r}) = s(L_{1,r}) + 1$ for $r \in \{5, 7, 9, 11, 13\}$. It follows that $L_{1,r} = T(2, r)$ is possible. However, $|\text{Kh}(L_{1,r})| = 3$ for such r , so they cannot be alternating knots.

When r is even, we note that $L_{q,r} = U \cup J_q$, where J_q is the 2-bridge knot $K[2/(2q - 9)]$. This knot is only a torus knot (or the unknot) if $q = 3, 4, 5$, or 6 , so $L_{q,r}$ can only be a Seifert link for these values, by Criterion 2.19. However, $L_{4,r}$ is never alternating, so it cannot be $T(2, 2k)$, and, $L_{3,r}$ has at least 9 crossings, so it cannot be the trefoil union a core curve.

We show $L_{q,r}$ cannot be a Montesinos knot or link by applying Method 1. When r is even, we note that one tangle would be a $(2/(2q + 5))$ -tangle or a $((q + 3)/(2q + 5))$ -tangle.

REFERENCES

- [AP04] Marta Asaeda and Józef Przytycki, *Khovanov homology: torsion and thickness*, Advances in topological quantum field theory, no. 179 of NATO Scientific Series II: Mathematics, Physics and Chemistry, Kluwer Academic Publishers, Dordrecht, 2004.
- [BGZ11] Steven Boyer, Cameron McA. Gordon, and Xingru Zhang, *Dehn fillings of knot manifolds containing essential once-punctured tori*, arXiv:1109.5151v3 [math.GT] (2011).
- [BGZ12] ———, *Characteristic submanifold theory and toroidal Dehn filling*, Advances in Mathematics **230** (2012), no. 4-6, 1673–1737.
- [BM70] Gerhard Burde and Kunio Murasugi, *Links and Seifert fiber spaces*, Duke Mathematics Journal **37** (1970), 89–93.
- [Bri98] Mark Brittenham, *Exceptional Seifert-fibered spaces and Dehn surgery on 2-bridge knots*, Topology **37** (1998), no. 3, 665–672.
- [BW01] Mark Brittenham and Ying-Qing Wu, *The classification of exceptional Dehn surgeries on 2-bridge knots*, Communications in Analysis and Geometry **9** (2001), no. 1, 97–113.
- [CGLS87] Marc Culler, Cameron McA. Gordon, John Luecke, and Peter Shalen, *Dehn surgery on knots*, Annals of Mathematics (2) **125** (1987), no. 2, 237–300.
- [Con70] John H. Conway, *An enumeration of knots and links, and some of their algebraic properties*, Computational Problems in Abstract Algebra (Pergamon, Oxford) (J. Leech, ed.), vol. Proc. Conf., Oxford, 1967, 1970, pp. 329–358.
- [Cro04] Peter R. Cromwell, *Knots and links*, Cambridge University Press, Cambridge, 2004.
- [Deh10] Max Dehn, *Über die Topologie des dreidimensionalen Raumes*, Mathematische Annalen **69** (1910), no. 1, 137–168.
- [Del95] Charles Delman, *Essential laminations and Dehn surgery on 2-bridge knots*, Topology and its Applications **63** (1995), no. 3, 201–221.
- [EM97] Mario Eudave-Muñoz, *Non-hyperbolic manifolds obtained by Dehn surgery on hyperbolic knots*, Geometric Topology, Studies in Advanced Mathematics (William H. Kazez, ed.), vol. 2.1, AMS and International Press, 1997, pp. 35–61.
- [EN85] David Eisenbud and Walter Neumann, *Three-dimensional link theory and invariants of plane curve singularities*, Annals of Mathematics Studies, 110, Princeton University Press, Princeton, NJ, 1985.
- [FIK⁺09] David Futer, Masaharu Ishikawa, Yuichi Kabaya, Thomas Mattman, and Koya Shimokawa, *Finite surgeries on three-tangle pretzel knots*, Algebraic & Geometric Topology **9** (2009), no. 2, 743–771.
- [Gab89] David Gabai, *Surgery on knots in solid tori*, Topology **28** (1989), no. 1, 1–6.
- [GL89] Cameron McA. Gordon and John Luecke, *Knots are determined by their complements*, Journal of the American Mathematical Society **2** (1989), no. 2, 371–415.
- [Gor99] Cameron McA. Gordon, *Small surfaces and Dehn filling*, Proceedings of the Kirbyfest (Berkeley, CA 1998) (Coventry), vol. 2, Geometry and Topology Monographs, Geometry and Topology Publishing, 1999, pp. 177–199.
- [Gor09] ———, *Dehn surgery and 3-manifolds*, Low dimensional topology **IAS/Park City Mathematics Series**, **15** (2009), 21–71.
- [Hei74] Wolfgang Heil, *Elementary surgery on Seifert fiber spaces*, Yokohama Math Journal **22** (1974), 135–139.
- [IJ09] Kazuhiro Ichihara and In Dae Jong, *Cyclic and finite surgeries on Montesinos knots*, Algebraic & Geometric Topology **9** (2009), no. 2, 731–742.
- [IJ11] ———, *Seifert fibered surgery and Rasmussen invariant*, arXiv:1102.1117v1 [math.GT], 2011.
- [IJK11] Kazuhiro Ichihara, In Dae Jong, and Yuichi Kabaya, *Exceptional surgeries on $(-2, p, p)$ -pretzel knots*, arXiv:1102.1118v1 [math.GT], 2011.
- [Kan10] Sungmo Kang, *Examples of reducible and finite Dehn fillings*, Journal of Knot Theory and its Ramifications **19** (2010), no. 5, 677–694.
- [Lee11] Sangyop Lee, *Lens spaces and toroidal Dehn fillings*, Mathematische Zeitschrift **267** (2011), no. 3-4, 781–802.
- [LM08] Marc Lackenby and Robert Meyerhoff, *The maximal number of exceptional Dehn surgeries*, arXiv:0808.1176v1 [math.GT] (2008).

- [LT88] W. B. Raymond Lickorish and Morwen B. Thistlethwaite, *Some links with nontrivial polynomials and their crossing-numbers*, Commentarii Mathematici Helvetici **63** (1988), no. 4, 527–539.
- [MB84] John W. Morgan and Hyman Bass (eds.), *The Smith Conjecture*, Pure and Applied Mathematics, 112, Academic Press, Orlando, FL, 1984.
- [MM02] Katura Miyazaki and Kimihiko Motegi, *Seifert fibering surgery on periodic knots*, Topology and its Applications **121** (2002), no. 1-2, 275–285.
- [Mon75] José M. Montesinos, *Surgery on links and double branched covers of S^3* , Knots, groups, and 3-manifolds (Papers dedicated to the memory of R. H. Fox, vol. 84, Princeton University Press, 1975, pp. 227–259.
- [Mot03] Kimihiko Motegi, *Dehn surgeries, group actions, and Seifert fiber spaces*, Communications in Analysis and Geometry **11** (2003), no. 2, 343–389.
- [Oer84] Ulrich Oertel, *Closed incompressible surfaces in complements of star links*, Pacific Journal of Mathematics **111** (1984), no. 1, 209–230.
- [Per03a] Grisha Perelman, *The entropy formula for Ricci flow and its geometric applications*, arXiv:math.DG/0211159 (2003).
- [Per03b] ———, *Finite extinction time for the solutions to the Ricci flow on certain three-manifolds*, arXiv:math.DG/0307245 (2003).
- [Per03c] ———, *Ricci flow with surgery on three-manifolds*, arXiv:math.DG/0303109 (2003).
- [Ras10] Jacob Rasmussen, *Khovanov homology and the slice genus*, Inventiones Mathematicae **182** (2010), no. 2, 419–447.
- [Wol99] Stephan Wolfram, *The Mathematica® Book*, fourth ed., Wolfram Media, Inc., Champaign, IL; Cambridge University Press, Cambridge, 1999.
- [Wu96] Ying-Qing Wu, *Dehn surgery on arborescent knots*, Journal of Differential Geometry **43** (1996), no. 1, 171–197.
- [Wu98a] ———, *Dehn surgery on arborescent knots and links-a survey*, Chaos Solitons Fractals **9** (1998), no. 4-5, 671–679.
- [Wu98b] ———, *Sutured manifold hierarchies, essential laminations, and Dehn surgery*, Journal of Differential Geometry **48** (1998), no. 3, 407–437.
- [Wu10] ———, *Persistently laminar branched surfaces*, arXiv:1008.2680v1 [math.GT], 2010.
- [Wu11a] ———, *The classification of toroidal Dehn surgeries on Montesinos knots*, Communications in Analysis and Geometry **19** (2011), no. 2, 305–345.
- [Wu11b] ———, *Exceptional Dehn surgery on large arborescent knots*, Pacific Journal of Mathematics **252** (2011), no. 1, 219–243.
- [Wu11c] ———, *Immersed surfaces and Seifert fibered surgery on Montesinos knots*, arXiv:0910.4882v2 [math.GT], 2011.
- [Wu12a] ———, *Dehn surgery on knots of wrapping number 2*, arXiv:1105.4287v1, 2012.
- [Wu12b] ———, *Seifert fibered surgery on Montesinos knots*, arXiv:12007.0154v1, 2012.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF TEXAS AT AUSTIN, 1 UNIVERSITY STATION, AUSTIN, TX 78712

E-mail address: jmeier@math.utexas.edu